


**REDOX INTERACTIONS BETWEEN IRON AND CARBON IN PLANETARY MANTLES: IMPLICATIONS FOR DEGASSING AND MELTING PROCESSES.** A. Martin and K. Righter, NASA Johnson Space Center, 2101 NASA Parkway, Houston, TX 77058, USA

**Introduction:** Carbon stability in planetary mantles has been studied by numerous authors because it is thought to be the source of C-bearing atmospheres and of C-rich lavas observed at the planetary surface. In the Earth, carbonaceous peridotites and eclogites compositions have been experimentally studied at mantle conditions [1] [2] [3]. [4] showed that the  $f_{O_2}$  variations observed in martian meteorites can be explained by polybaric graphite-CO-CO<sub>2</sub> equilibria in the Martian mantle. Based on thermodynamic calculations [4] and [5] inferred that the stable form of carbon in the source regions of the Martian basalts should be graphite (and/or diamond), and equilibrium with melts would be a source of CO<sub>2</sub> for the martian atmosphere. Considering the high content of iron in the Martian mantle (~18.0 wt% FeO; [6]), compared to Earth's mantle (8.0 wt% FeO; [7]) Fe/C redox interactions should be studied in more detail.


**Experiments:** Here we report new experimental results at 2 GPa (piston-cylinder) in a system containing CaO-MgO-SiO<sub>2</sub>-CO<sub>2</sub> with various FeO content and oxidation conditions. By analyzing Fe in liquid and solid phases, we have determined Fe repartition and constrained Fe/C redox interactions during late differentiation processes in terrestrial planets.

**Results and discussion:** In the Fe-free system from 1100 to 1300°C, CO<sub>2</sub>-rich melt forms in equilibrium with forsterite, clinopyroxene and CO<sub>2</sub>. In the presence of Fe, forsterite disappears and Fe-opx is produced. Fe oxide (magnetite) is also found and part of the carbon is reduced to graphite (Fig.1).



**Figure 1:** SEM image (back-scattered electrons) of a 2.0 GPa - 1200°C sample containing Fe-opx (light grey), Mg-cpx (medium grey), calcite (dark grey), CO<sub>2</sub>-rich melt (interstitial), graphite (black) and Fe-oxides (white).

Thus, a reaction occurs between iron- and carbon-bearing phases:



Such supersolidus redox process will have an influence on the melting and degassing mechanisms. Especially, it may lower the contribution of late mantle degassing processes to CO<sub>2</sub> introduction into planetary atmospheres.

**References:** [1] Dasgupta R. and Hirschmann M.M. (2007) *American Mineralogist*, 92, 370–379. [2] Brey G.P. et al. (2008) *Journal of Petrology*, 49, 4, 797-821. [3] Hammouda T. (2003) *EPSL*, 214, 1-2, 357-368. [4] Righter K. et al. (2008) *Meteoritics and Planet. Sci.*, 43, 10, 1709-1723. [5] Hirschmann M.M. and Withers A.C. (2008) *EPSL*, 270, 147-155. [6] Dreibus and Wänke (1985) *Meteoritics*, 20, 367-382. [7] McDonough W. F. and Sun S.-s. (1995) *Chemical Geology*, 120, 223-254.